

Natural Color Transparency in High Energy (p,pp) Reactions

B.K. Jennings

TRIUMF, 4004 Wesbrook Mall, Vancouver, B.C., V6T 2A3, Canada

and

G.A. Miller

Department of Physics, FM-15, University of Washington, Seattle, WA 98195, USA

ABSTRACT

New parameter free calculations including a variety of necessary kinematic and dynamic effects show that the results of BNL ($p, 2p$) measurements are consistent with the expectations of color transparency.

(submitted to Phys. Lett. B)

The anomalously large transmission of a hadron in the nuclear medium following or preceding a hard interaction is commonly referred to as color transparency[1, 2]. This topic has been actively studied theoretically [3]-[9]. Color transparency depends on the formation of a small-sized wavepacket or point-like configuration PLC in a high momentum transfer quasi-elastic reaction. The effects of color screening or color neutrality [10] suppress the interaction between the PLC and the nuclear medium. At the present experimental energies the PLC expands as it moves through the nucleus and therefore does interact. This expansion is treated in Refs.[3, 8, 9, 11, 12].

Color transparency is under active experimental investigation at BNL [13, 14] and at SLAC[15, 16]. The $(p, 2p)$ experiment of Carroll et al[13] finds a transparency T (ratio of nuclear to hydrogen cross section per nucleon after removing the effects of nucleon motion) with an oscillatory pattern. T increases as the beam momentum increases from 5 to 10 GeV/c but then decreases. The SLAC experiment is unpublished at this time, but a preliminary report [16] found no effect of color transparency (CT) in $(e, e'p)$ reactions for Q^2 between 1 and 7 GeV^2 . However, the stated systematic errors are about $\pm 10\%$ which are large compared to the size of the predicted CT effects.

The purpose of this note is to include all of the known necessary kinematic and dynamic effects: a proper treatment of the longitudinal (parallel to the virtual photon or incident proton momentum) component of the momentum of the detected protons[17, 18]; computing the measured experimental observable which involves an integration over the transverse momentum of the struck proton; including the effects on interference between PLC and non-PLC configurations produced in proton-proton elastic scattering [6, 7]; and, a realistic treatment of the baryonic components of the PLC[12]. Thus our aim here is to show that including the three kinematic and dynamic effects mentioned above along with the color transparency of ref. [12] leads to a natural explanation of the existing BNL data. No free parameters are used in the present work.

We start the analysis by recalling the mechanisms proposed to understand the (p, pp) reaction. The suggestion of Ralston and Pire [7], is that the energy dependence is caused

by an interference between a hard amplitude which produces a PLC, and a soft one (blob-like configuration BLC) which does not. The Ralston and Pire idea is that the BLC is due to the Landshoff process[19]. Another mechanism with a similar effect is that of Brodsky and de Teramond [6] in which the two-baryon system couples to charmed quarks (there is a small (6q) and a BLC which is a (6q, $c\bar{c}$) object) The Brodsky-de Teramond idea is motivated by the fact that the mass scale of the rapid energy variation in A_{NN} [20] and in the measured transparency matches that of the charm threshold.

Both of the mechanisms of [7] and [6] are well motivated, but imprecisely understood. For example, the nature of the Landshoff term is uncertain. At high energies one expects that this component is suppressed by a Sudakov factor [21]. At present energies, the energy dependence and phase of the term are not well determined, and the size of the object produced by the Landshoff process is not well known. See e.g. ref. [22]. Nevertheless, the Sudakov effects can be expected to a set of configurations with a range of different sizes. Similarly even if the details of the $c\bar{c}$ production amplitudes are not yet well-established, threshold effects will naturally lead to mixtures of BLC and PLC. Thus, it's natural to discuss high Q^2 elastic proton-proton scattering in terms of configurations of different sizes. Separating the contributing configurations into two, a PLC and a BLC is only a simple first step.

The two ideas about the BLC can be used to qualitatively explain the oscillations observed in the (p,pp) data, but do not quantitatively reproduce features of the data Ref. [23, 11] when combined with a proper treatment of the expansion[11]. The agreement is not quantitative because including the non-zero absorption of the expanding PLC and the non-complete absorption of the BLC tends to make the two terms similar and weakens the interference effects.

Another relevant effect is to properly account for the correct momentum of the detected protons. Consider for example, the (e,e'p) reaction in which the virtual photon has the momentum $\vec{q}$. Suppose the detected proton has momentum $\vec{p}$ with $\vec{p} = \vec{q} + \vec{k}$. For quasi-elastic kinematics $\vec{p} = \vec{q}$ and $\vec{k} = \vec{0}$. Fermi motion or final state interactions

can lead to a non-zero value of $\vec{k}$. This is old news. However the non-zero components of $\vec{k}$ in the direction of $\vec{q}$ (k_z) have been shown to give a large numerical effect [17]. A detailed study of this “Fermi motion” effect was made by Frankfurt et al [18]. The momentum $\vec{k}$ is sometimes called the momentum of the struck nucleon (and is that quantity if the plane wave Born approximation is valid). We also use the term “momentum of the struck proton”, but only as an abbreviation. For the (p,pp) reaction the momentum of the detected protons can also be described in terms of the momentum of the struck nucleon [13]. Indeed, the data are presented in terms of bins of k_z , the component of $\vec{k}$ parallel to the beam direction. This is especially important for the (p,pp) reaction because the proton- proton scattering cross section varies approximately as s^{-10} and here $s = 2M^2 + 2ME_{LAB} - 2k_z P_{LAB} \approx 2MP_{LAB}(1 - k_z/M)$ where M is the proton mass. Ref [18] examined the (p,pp) reaction, including the effects of non-zero values of k_z in their calculation, and obtained qualitative agreement with the 10 GeV data, but not with data taken at the other energies. Agreement was also obtained by Kopeliovich[24] for some energies. Neither group included any effects of interference between PLC and BLC.

Our calculation of the nuclear (p,pp) cross sections uses the quantum mechanical treatment of color transparency developed in ref. [8, 9, 12]. In that approach the time for the PLC to expand depends on the masses of the baryonic components of the PLC. In early work [8, 9] only a single average state of mass M_1 was used. Then the time for the PLC to expand is $\frac{2P_{LAB}}{(M+M_1)} \frac{1}{(M_1-M)}$. Recently we [12] included the effects of the continuous baryon spectrum by using measured proton diffractive dissociation and electron deep inelastic scattering data to constrain the baryon masses. Our discussion of the formalism shall be brief and concentrate on the newer kinematic and dynamic aspects. For details see Refs. [9, 11, 12].

To be definite, consider the (e,e'p) high momentum transfer process in which a photon of three-momentum $\vec{q}$ is absorbed and a nucleon of momentum $\vec{p}$ leaves the nucleus. As usual, $q^2 = -Q^2$. Consider knockout from only a single shell model orbital, denoted by α . The observable cross sections are computed by making incoherent sums over all the

occupied orbitals.

Let the amplitude be defined as $\mathcal{M}_\alpha$, which can be described in terms of a series involving multiple wavepacket (PLC) -nucleon scatterings. The first term is the plane wave Born term B_α in which the wavepacket undergoes no interactions. If full color transparency is obtained, this is the only term to survive. The first correction to the Born term is the scattering term or second term denoted by ST_α . To the stated order we have: $\mathcal{M}_\alpha = B_\alpha + ST_\alpha$. The Born term is given by

$$B_\alpha = \langle \vec{p} | T_H(Q) | \alpha \rangle = F(Q^2) \langle \vec{p} - \vec{q} | \alpha \rangle, \quad (1)$$

in which $T_H(Q^2)$ is the hard scattering operator. Specific effects of spin are ignored here and throughout this work.

The second term is denoted by

$$ST_\alpha = \langle \vec{p} | U G T_H(Q) | \alpha \rangle, \quad (2)$$

where G is the Green's operator for the emerging small object (PLC). This object is a wave packet of which the nucleon is just one component. The operator U represents the interaction between the ejected baryon and the nuclear medium. The nuclear interaction can change the momentum of the ejectile and also excite or de-excite the internal degrees of freedom. Our approach is to treat the Green's function G as a sum of baryonic propagators each denoted by m . The eikonal Green's function for the emerging baryon is:

$$G(Z, Z') = \sum_m G_m(Z, Z') = \sum_m \frac{\theta(Z - Z') e^{ip_m(Z - Z')}}{2ip_m}, \quad (3)$$

with $\vec{p}_m^2 = \vec{p}^2 + M^2 - M_m^2$. Here M is the nucleon mass, M_m the mass of the baryonic state. Our notation is that $p = |\vec{p}|$ and $p_m = |\vec{p}_m|$. Terms beyond the first order in U are included by exponentiation, which is an excellent approximation for our applications [25].

The work of ref. [9] involves a simple model in which the operator U gives only one

excited state (m=1) when acting on a nucleon. In that model

$$ST_\alpha = -F(Q^2) \int d^2 B dZ \rho(B, Z) \frac{e^{-ipZ}}{(2\pi)^{3/2}} \int_{-\infty}^Z dZ' e^{-ip(Z'-Z)} \frac{\sigma_{eff}}{2}(Z, Z') e^{iqZ'} \langle \vec{B}, Z' | \alpha \rangle, \quad (4)$$

where the $\rho(B, Z)$ is the nuclear density, and

$$\sigma_{eff}(Z, Z') \equiv \sigma \left(1 - e^{i(p-p_1)(Z'-Z)} \right), \quad (5)$$

and σ is the proton-nucleon total cross section. This is eq (32) of ref. [9].

Now we can exhibit the importance of the momentum (k_z) of the struck nucleon. We take $p = q + k_z$ for components parallel to the direction of $\vec{q}$. Change the integration variable Z' to D via $Z' = Z - D$. Then

$$ST_\alpha = -F(Q^2) \int d^2 B dZ \sigma \rho(B, Z) \frac{e^{-ik_z Z}}{(2\pi)^{3/2}} \int_0^\infty dD e^{ik_z D} \left(1 - e^{-i(p-p_1)D} \right) \langle \vec{B}, Z' | \alpha \rangle. \quad (6)$$

The effect of k_z appears in two places. The $e^{-ik_z Z}$ factor is the same as in standard Born or Distorted Wave Born calculations (DWBA or Glauber optical model). The $e^{ik_z D}$ factor involves a modification of the color transparency physics. To see this note that the real part of ST_α dominates the numerics. Then $\cos(k_z D) - \cos((k_z - p + p_1)D) \approx \frac{D^2}{2}(p - p_1)(p - p_1 - 2k_z)$ for small D . Since $p > p_1$ a positive value of k_z reduces this term and increases the transparency for any given value of p . This latter effect does not occur in the color transparency of ref. [3], but would occur in models in which the baryon-nucleon interaction is treated as a finite-dimensional matrix[24, 26].

The above paragraph is meant as a simple explanation of a numerical effect that is surprisingly large. The computations of (p, pp) reactions are more involved since there is one incident proton wave function and two outgoing ones. Furthermore the amplitudes for the production of both the PLC and BLC must be taken into account. However the qualitative effect of including non-zero values of k_z (here z is the direction of the beam proton) is similar to the electron scattering case. Note also that our more realistic calculations [12] replace the single mass M_1 by an appropriate distribution of masses.

The next step is to discuss the observables measured in ref.[13]. Let the four-momentum of the target proton be denoted as $(M, \vec{k})$. The transparency T is defined as a ratio $T = d\sigma/d\sigma^B$ with

$$d\sigma = \int_{k_a}^{k_b} dk_z \int \int dk_x dk_y (d\sigma^A/dt)(s) \quad (7)$$

and

$$d\sigma^B = \int_{k_a}^{k_b} dk_z \int \int dk_x dk_y (d\sigma^B/dt)(s) \quad (8)$$

in which the superscript A denotes the nuclear cross section divided by the number of target protons and the subscript B denotes the same quantity but computed in the plane wave Born approximation. In the experiment the integrations over the transverse momenta k_x, k_y were limited to about 250 MeV/c. This corresponds to almost all of the probability so we integrate over all k_x, k_y in our calculations. The integration over the transverse components of $\vec{k}$ reduces the Glauber DWBA result for T by about 30%. The data of ref. [13] are presented in terms of T for each (k_a, k_b) .

We now present the results. The calculations for an Al target (three beam momenta and four bins of k_z) are shown in Fig 1. The solid curves show the effect of CT using the complete power law form of the distribution of baryon masses as in ref. [12] and dashed curves show the results of using Ref.[9] with a value of $M_1 = 1550 \text{ MeV}$. This value gives small enough color transparency at low Q^2 so as to be consistent with the NE-18 data. The Ralston-Pire parameterization of $d\sigma/dt$ for the free protons is used here along with their separation of the BLC and PLC terms. Details of the implementation of this are to be found in ref [11]. The use of the corresponding Brodsky-de Teramond model for the PLC-BLC interference would lead to similar results for the energy range we consider here [11, 12]. The data are from ref. [13]. The target-dependent and target independent uncertainties in the normalization of T are about 10% and 25% respectively [13]. We multiply the central values of the Al, Cu, and Pb data points by a factor of 0.75, and those of Li and C by 0.85. This is consistent with the published errors. Each data “point” represents a bin of k_z , represented as a horizontal line. The integration of Eqs. (7),(8) over the small bins 0.1 or 0.2 GeV/c of ref.[13] causes negligible differences with simply using the central values. The use of a distribution of masses starting at $M + m_\pi$ increases the computed $d\sigma/d\sigma_B$ at the lowest beam momentum. In either case, including

the effects of color transparency gives good agreement with the data.

Fig. 2, shows a summary of the different calculations for ^{27}Al . Here as in Figs. 1 and 3 the solid curve represents the full color transparency calculation, the dashed curves are obtained by neglecting the effects of the PLC-BLC interference, and the dotted curves represent the use of the standard optical model or “Glauber” calculation. These latter curves fall far below the data, but including the effects of CT leads to a reasonable reproduction of the data.

Fig. 3 shows the A-dependence of the data of ref.[13]. These are taken for a bin of k_z ranging from -0.2 GeV/c to 0.1 GeV/c. Here the use of Eqs. (7) and (8) does matter. To see this compare the ^{27}Al data of this figure with the previous figures. The Glauber standard optical model leads to results (dotted curves) that again fall well below the data, this time for each target nucleus. The solid curves, which show our full calculation, are in excellent agreement with all of the data, except for the 12 GeV Al data point. The dashed curves show the results obtained without the BLC-PLC interference Ralston-Pire effect, so we see that the latter helps to account for the energy dependence, even though it is not a very large numerical effect.

It is necessary to comment on the single particle nuclear shell model wave functions used here. Harmonic oscillator ($\hbar\omega = 41\text{MeV}/A^{1/3}$) wave functions are used for light nuclei (Li and C). The other nuclei are treated in the Hartree-Fock (HF) approximation with the SGII interaction of Ref. [27]. (HF wave functions are used in obtaining the results of Figs. 1,2.) If the oscillator frequency is chosen appropriately, the HF wave functions are well approximated by a single harmonic oscillator wave function. Thus we find the most important effect of using the HF wavefunctions is to shift the value of $\hbar\omega$ from $41\text{MeV}/A^{1/3}$ to one that more precisely represents the nuclear mean square radius. The effect of using HF wave functions with the correct exponential dependence at long distance on $d\sigma/d\sigma^B$ is largest, a 20% increase independent of energy, for the Glauber optical model calculations for the Pb target. But the effect is negligible for the CT calculations. See also ref.[28].

The main effect of the color transparency is to provide an increase in the predicted magnitude of the cross sections. One might wonder if some combination of reasonable effects applied to the Glauber optical model calculations yield enhancements that lead to reproducing the data without including color transparency effects. One possibility is to claim [29] that the experimental resolution allows the inclusion of all nuclear excited states in the measurement. In that case, the optical model wave functions would be computed using the proton-proton reaction cross section σ_r instead of the total cross section σ as we have done. However, $\sigma_r/\sigma \approx 0.85$ [30] and has only a small energy dependence, for the energy range of interest here. Calculations show that including this effect provides only 10-20 % enhancements. Including the effects of nucleon-nucleon short range correlations leads to a 20% increase for ^{12}C [28], but is a much smaller effect for heavier nuclei. Thus, in the absence of CT effects, we see no possibility to elevate the DWBA cross sections to the levels observed by the experiment.

We believe the BNL experiment and the present calculations calibrate the size of color transparency effects. Thus we determine the energies for measurable effects. Our calculations provide a good guide to the (e,e'p) experiments. The SLAC Ne-18 experiment is set for very small values of k_z , so the previous results of ref [12] stand unchanged (except possibly for the effects of using improved nuclear structure information for heavy targets). Since the predicted transparency is not large in the range of Q^2 between 1 and 7 GeV^2 , we expect that our earlier calculations of color transparency effects for the power-law form of $g(M_X^2)$ will not be ruled out by the final results of the SLAC experiment [15, 16]. Furthermore, our CT calculations predict enhancements for Q^2 between 7 and 15 GeV^2 . Thus a higher energy experiment which examines the k_z dependence should observe measurable effects.

Thus, we summarize: If CT effects are included, the qualitative features of the data of ref [13] can be reproduced in a very natural way. No adjustment of parameters is needed. If the new BNL (p,pp) color transparency experiment [14] confirms the central values of the older results, one can be confident that color transparency has been discovered.

Acknowledgment: One of the authors (BKJ) thanks the Natural Sciences and Engineering Research Council of Canada for financial support. The other (GAM) thanks the DOE for partial support. Discussions with W.R. Greenberg and M. Strikman are appreciated. We thank G. Bertsch for making the HF program of ref. [27] accessible to us.

References

- [1] A.H. Mueller in Proceedings of Seventeenth rencontre de Moriond, Moriond, 1982 ed. J Tran Thanh Van (Editions Frontieres, Gif-sur-Yvette, France, 1982)p13.
- [2] S.J. Brodsky in Proceedings of the Thirteenth intl Symposium on Multiparticle Dynamics, ed. W. Kittel, W. Metzger and A. Stergiou (World Scientific, Singapore 1982,) p963.
- [3] G.R. Farrar, H. Liu, L.L. Frankfurt & M.I. Strikman, Phys. Rev. Lett. 61 (1988) 686.
- [4] L. Frankfurt and M. Strikman, Prog. Part. and Nucl. Phys., 27,135,(1991); L. Frankfurt and M. Strikman, Phys. Rep. 160, 235 (1988).
- [5] L Frankfurt, G.A. Miller & M. Strikman, Comm. Nuc. Part. Phys. 21,1 (1992).
- [6] S.J. Brodsky & G.F. De Teramond, Phys. Rev. Lett. 60 (1988) 1924.
- [7] J.P. Ralston and B. Pire, Phys. Rev. Lett. 61 (1988) 1823
- [8] B.K. Jennings and G.A. Miller, Phys. Lett. B236 (1990) 209.
- [9] B.K. Jennings and G.A. Miller, Phys. Rev. D 44,692 (1991)
- [10] S. Nussinov Phys. Rev. Lett 34, 1286 (1975); F.E. Low, Phys. Rev. D12, 163 (1975); J. Gunion D. Soper ,*ibid* D15, 2617 (1977).

- [11] B.K. Jennings and G.A. Miller Phys. Lett. B274,442 (1992).
- [12] B.K. Jennings and G.A. Miller Phys. Rev. Lett. 69,3619 (1992).
- [13] A.S. Carroll et al. Phys. Rev. Lett. 61 (1988) 1698; S. Heppelmann, p.199 in “Nuclear Physics and the Light Cone”, ed. by M.B. Johnson and L.S. Kisslinger, World Scientific (Singapore, 1989)
- [14] A.S. Carroll, S. Heppelmann et al, BNL experiment 850 EVA, 1991.
- [15] R. Milner et al., SLAC proposal NE18 and private communications.
- [16] R.D. McKeown Bull. Amer. Phys. Soc. 37 (1992) 1254.
- [17] B Kopeliovich and B K Jennings 1992 TRIUMF preprint
- [18] L Frankfurt, M. Strikman, and M. Zhalov 1993 PSU preprint
- [19] B. Pire and J.P. Ralston, Phys. Lett. 117B (1982) 117B, J.P Ralston and B. Pire, Phys. Rev. Lett. 49 (1982) 1605.
- [20] E. A. Crosbie et al. Phys Rev. D23,600(1981)
- [21] A. H. Mueller, p. 185 in ”Nuclear Physics on the Light Cone”, Ed. by M. B. Johnson and L.S.Kisslinger, World Scientific, Singapore 1988.
- [22] J. Botts and G. Sterman, Nucl. Phys. B325,62(1989); Phys.Lett.B224,201(1989),E-ibid.B227,501(1989). J. Botts, Phys.Rev.D44,2768 (1991); University Park Workshop 1990,230; Nucl.Phys.B353,20(1991). J. Botts, J-W Qiu, G. Sterman, Nucl.Phys.A527,577 (1991). Phys.Lett.B224,201(1989),E-ibid.B227,501(1989).
- [23] B.Z. Kopeliovich and B.G. Zakharov, Phys. Lett. B264 (1991) 434.
- [24] B.Z. Kopeliovich, private communication.

- [25] The exponentiation procedure is also a very accurate approximation to the appropriate multiple-scattering series, W. R. Greenberg and G.A. Miller, Phys. Rev. D47,1865 (1993). This is equivalent to replacing σ by σ_{eff} (see Eqs. 45) instead of in standard scattering wave functions. σ_{eff} is complex because it is part of a contribution to a scattering amplitude.
- [26] L Frankfurt, W.R. Greenberg, G.A. Miller, and M. Strikman Phys. Rev. C46, 2547 (1992).
- [27] N. Van Giai and H. Sagawa, Phys. Lett. 106B (1979) 379.
- [28] T.-S.H. Lee and G.A. Miller, Phys. Rev C. 45, 1863 (1992).
- [29] A. Kohama, K. Yazaki and, R. Seki, Nucl. Phys. A536 (1992) 716.
- [30] PDG, Phys. Rev D 45, Number 11, Part II, 1992

Figure Captions

Figure 1: Effect of using distributed baryonic masses. Solid uses distributed masses. Dashed- $M_1 = 1550$ MeV. The data in all figures are those of Ref. [13].

Figure 2. Full Al data. Solid- full calculation. Dashed- with out the Ralston-Pire effect. Dotted -Glauber.

Figure 3. A dependence of transparency. Solid- full calculation. Dashed- with out the Ralston-Pire effect. Dotted -Glauber.

This figure "fig2-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9305317v1>

This figure "fig2-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9305317v1>

This figure "fig2-3.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9305317v1>

This figure "fig2-4.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9305317v1>